



Shakespeare, C., Merriel, A., Bakhbakhi, D., Baneszova, R., Barnard, K., Lynch, M., Storey, C., Blencowe, H., Boyle, F., Flenady, V., Gold, K., Horey, D., Mills, T., & Siassakos, D. (2019). Parents' and healthcare professionals' experiences of care after stillbirth in low- and middle-income countries: a systematic review and meta-summary. *BJOG: An International Journal of Obstetrics and Gynaecology*, 126(1), 12-21. <https://doi.org/10.1111/1471-0528.15430>

Peer reviewed version

Link to published version (if available):  
[10.1111/1471-0528.15430](https://doi.org/10.1111/1471-0528.15430)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://obgyn.onlinelibrary.wiley.com/doi/abs/10.1111/1471-0528.15430> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# Parents' and healthcare professionals' experiences of care after stillbirth in low and middle-income countries: a systematic review and metasummary'

Clare Shakespeare<sup>1</sup>, Abi Merriel<sup>1</sup>, Danya Bakhbakhi<sup>1</sup>, Ruth Banaszova<sup>2</sup>, Katie Barnard<sup>3</sup>, Mary Lynch<sup>4</sup>, Claire Storey<sup>5</sup>, Hannah Blencowe<sup>6</sup>, Fran Boyle<sup>7</sup>, Vicki Flenady<sup>7</sup>, Katherine Gold<sup>8</sup>, Dell Horey<sup>9</sup>, Tracey Mills<sup>10</sup>, Dimitrios Siassakos<sup>1</sup>

1. School of Population Health sciences, Bristol Medical School, University of Bristol, Department of Women and Children's Health, The Chilterns, Southmead Hospital, Bristol BS10 5NB.

2. 2<sup>nd</sup> Department of Obstetrics and Gynaecology, University Hospital Bratislava and Faculty of Medicine, Comenius University, Bratislava, Slovakia.

3. Library and Knowledge Service, Southmead Hospital, North Bristol NHS Trust, BS10 5NB.

4. Department of Women and Children's Health, The Chilterns, Southmead Hospital, Bristol BS10 5NB.

5. International Stillbirth Alliance, 61 Kings Drive, Stoke Gifford, Bristol, BS34 8RD. 6. London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT.

7. Centre of Research Excellence in Stillbirth, Mater Research Institute, University of Queensland, South Brisbane, Australia.

8. Department of Medicine, Department of Obstetrics and Gynaecology, University of Michigan, Ann Arbor, MI.

9. La Trobe University, Kingsbury Drive, Bundoora, Victoria 3086, Australia

10. School of Nursing, Midwifery and Social Work. University of Manchester, M13 9PL

Corresponding author: Dr Clare Shakespeare, The Chilterns, Southmead Hospital, Bristol, BS10 5NB. Phone: 07717798582. Email: [clare.shakespeare@gmail.com](mailto:clare.shakespeare@gmail.com)

Shortened running title (60 characters): Care after stillbirth in low and middle-income countries

Appendix S1: Search strategy

Appendix S2: Data extraction form

Appendix S 3: Summary of studies

Appendix S 4: Sample extracted findings

Appendix S 5: Intensity effect sizes

Appendix S 6: Quality assessment

## **Abstract:**

**Background:** Stillbirth has a profound impact on women, families, and healthcare workers. The burden is highest in low and middle-income countries (LMICs). There is need for respectful and supportive care for women, partners, and families after bereavement.

**Objective:** To perform a qualitative metasummary of parents' and healthcare professionals' experiences of care after stillbirth in LMICs.

**Search Strategy:** Search terms were formulated by identifying all synonyms, thesaurus terms and variations for stillbirth. Databases searched were AMED, EMBASE, MEDLINE, PsychINFO, BNI, CINAHL.

**Selection Criteria:** Qualitative, quantitative, and mixed method studies that addressed parents' or healthcare professionals' experience of care after stillbirth in LMICs.

**Data Collection and Analysis:** Studies were screened and data extracted in duplicate. Data was analysed using the Sandelowski meta-summary technique that calculates frequency and intensity effect sizes (FES/IES).

**Main Results:** 118 full texts were screened and 34 studies from 17 countries were included. FES range was 15-68%. Most studies had IES 1.5-4.5. Women experience a broad range of manifestations of grief following stillbirth, which may not be recognised by healthcare workers, or in their communities. Lack of recognition exacerbates negative experiences of stigmatisation, blame, devaluation, and loss of social status. Adequately developed health systems, with trained and supported staff, are best equipped to provide the support and information that women want after stillbirth.

**Conclusions:** Basic interventions could have an immediate impact on the experiences of women and their families after stillbirth. Examples include public education to reduce stigma, promoting the respectful maternity care agenda, and investigating stillbirth appropriately.

**Funding:** N/A

**Keywords:** Stillbirth, LMIC, global health, bereavement care, systematic review, qualitative metasummary

## **Tweetable abstract:**

Reducing stigma, promoting respectful care and investigating stillbirth has positive impact after stillbirth for women and families in LMICs

## Introduction

Stillbirths hold an increasingly important place in the global maternal and newborn health agenda. The loss of a baby has a profound impact on women, families, communities and healthcare workers(1). The burden is highest in low and middle income countries(LMICs) where an estimated 98% of the 2.6 million stillbirths worldwide occurred in 2015(2).

Efforts to provide effective antenatal and intrapartum care are essential for improving maternal health and preventing stillbirth in LMICs(3). Effective bereavement care is vital for preventing negative short and long term outcomes for families.(1) The 2016 Lancet Ending Preventable Stillbirth Series called for a 'global consensus on a package of care after a death in pregnancy or childbirth... for the affected family, community and caregiver'(4).

A systematic review in 2016(5) addressed experiences of care after stillbirth by parents and healthcare professionals in high-income countries (HICs). Parental findings included the desire for support in memory making, and for increased public awareness and prioritisation of stillbirth(5). Staff behaviours and actions have a memorable impact on parents(5), underlining the need for improved training and care pathways to support staff. In the UK, this evidence has been synthesised and already used to inform national care pathways (*nbcpathway.org.uk*).

No previous literature synthesis has focussed on evidence from LMICs, where most stillbirths occur. There is a lack of evidence-based recommendations about care provided to women, partners and families who experience bereavement in LMICs (4,6). Difficulty in providing bereavement care due to lack of support or training is a source of stress and challenge for healthcare workers(1).

This systematic review identified studies from LMICs and assessed the available evidence to identify themes which are important to both parents and healthcare workers. The aim was to determine themes which could be used to inform training, guidelines, and a subsequent consensus on global bereavement care principles.

## Methods

### Objective

To systematically review and perform a qualitative metasummary of research surrounding parents' and healthcare professionals' experiences of care after stillbirth in LMICs.

### Search strategy

The search strategy (inception to May 2017) used text word variations and thesaurus terms for stillbirth, families, healthcare professionals, personal experience, and LMICs (Appendix S1). LMICs were defined by December 2016 World Bank classification(7). The databases searched were AMED, EMBASE, MEDLINE, PsychINFO, BNI, CINAHL. Conference abstracts from the International Stillbirth Alliance and First Candle conferences were hand-searched for eligibility. There were no date or language limits for the search. There was no patient and public involvement in the systematic review, and no funding was required. No relevant core outcome sets are available.

### Eligibility criteria

Qualitative, quantitative, and mixed method studies that addressed parents or healthcare professionals experience of care after stillbirth in LMICs were included. Many papers included a

combination of miscarriage, stillbirth and early neonatal death. In many LMICs it is difficult to accurately define gestational age at stillbirth, and definitions vary internationally. Therefore, to avoid losing useful and relevant data, only studies explicitly addressing miscarriage, fetal anomaly, and neonatal death alone were excluded. In studies with results divided by type of loss, findings specific to stillbirth were extracted.

No language restrictions were applied. Dissertations and conference abstracts were included but none contributed to the final review. Review articles, opinion pieces and books were excluded but hand searched for relevant references.

### **Study selection**

Two reviewers screened all abstracts using Covidence software(8), an online platform for systematic review collaboration. Disagreements were resolved by discussion with a third author.

### **Quality assessment**

Study quality and risk of bias was assessed using standard checklists for observational(9) and qualitative(10) studies. The overall quality assessment was synthesised using the Gough weight of evidence framework(11) to take into account the appropriateness of the study method as well as the relevance of each study to answer the review question. Studies were not excluded on basis of quality as the Sandelowski metasummary method allows for inclusion of all potentially useful findings.

### **Data extraction**

Data was independently extracted from included studies in duplicate by two authors (CSh, DB) using a standardised data extraction form (Appendix S2). The data extraction form was piloted before use and no changes made.

### **Data analysis**

Sandelowski's metasummary approach(12,13) was used for quantitative aggregation of qualitative and quantitative findings. Firstly, data was extracted from each paper, regarding the events or experiences investigated. Similar findings were then grouped into topics, enabling identification of recurring findings whilst preserving the breadth and complexity of the data. Concise but comprehensive thematic sentences were then derived. The core research team (CSh, DB, AM, RB, ML, CSt, DS) reviewed and discussed all the extracted findings and developed thematic sentences.

Finally, effect sizes were calculated to add a quantitative 'weight' to each finding and to each study(12,13). The frequency effect size (FES) reflects the relative magnitude of an abstracted finding within the included studies, by answering the question 'in how many studies does this finding appear?' ('number of studies with the finding' divided by 'total number of studies'). The intensity effect size (IES) reflects the impact of each study and how much it contributed to the final set of findings by defining the number of findings within it ('number of findings within the study' divided by 'total number of findings'). This helps identify findings only presented in weaker studies, as well as studies which contributed findings with a large FES.

## **Results**

### **Study selection**

Electronic searches revealed 2491 records, and an additional 13 were identified from handsearching conference abstracts and review references. After duplication and eligibility screening, 118 full texts were obtained. 34 were eligible for inclusion (Figure 1- Prisma diagram).

### **Description of included studies.**

Over half of the included studies used qualitative study designs (19, 56%), with quantitative (7, 21%) and mixed method (8, 24%) study designs playing a smaller role (Figure 2). A summary of the studies in the review is included as Appendix S3.

The 34 studies were conducted in 17 countries across 5 regions (Figure 2), by 26 different research groups. All but 5 were conducted in middle income countries (85%), with an equal number of studies in upper middle income countries (South Africa [5]; Brazil [3]; Iran [3]; Malaysia [2]; and one each in China [Hong Kong and Singapore findings excluded] and Russia) and in lower middle income countries (India [6 studies]; Nigeria [3]; Ghana [2]; and one each in Bangladesh, Indonesia and Uganda). Studies were conducted in the following low income countries: Benin; Ethiopia; Malawi; Somalia; Tanzania.

Overall the studies involved 2934 participants, including 1128 women who had experienced some form of perinatal loss; 300 were specifically identified as experiencing stillbirth. One study focussed on 33 men whose wife had had a stillbirth(14). Most studies (27, 79%) related to women's experiences of stillbirth or perinatal loss. Four studies included 259 health professionals with professional experience of stillbirth(15–18) .

### **Findings**

The analysis identified 372 individual 'findings' (Appendix S4). From these, 13 thematic sentences were agreed which are represented schematically in Figure 3. These were used for calculation of FES (Figure 4).

### **Thematic sentences**

***Positive community support, as opposed to stigmatisation and blame, can improve bereavement experience.***(14,19–40) (FES 68%): Women who experienced emotional and material support from society, including family, friends, religious and peer support groups reported lower perinatal grief and depression scores. Negative experiences on the other hand, included stigma, blame, loss of social status, social isolation, relationship difficulties, denial of motherhood and disenfranchised grief.

***Women's' experience of grief has multiple manifestations often unrecognised by the healthcare community and wider society.***(14,19,20,23–26,28–38,41–44) (FES 65%): Women across all cultures experienced grief that manifested as physical symptoms such as fatigue and pain, and emotionally, in the form of sadness, anxiety, guilt, confusion, and anger. Women wanted recognition of their baby, and of their own loss and bereavement. Fathers also experienced grief and sadness but this was often suppressed. The healthcare community, often failed to recognise and acknowledge grief, understand its context, or identify women's need for additional support.

***Awareness of, and support for, appropriate coping mechanisms can assist grieving .***(14,15,21–25,28–33,36–38,40–43,45,46) (FES 65%): A range of coping mechanisms that women found

beneficial were reported including religion and faith, distraction, family support, peer groups, self-medication, and traditional remedies. Other strategies, commonly used in HICs, were not universally supported, for example, taking pictures or seeing and holding the baby was not always culturally appropriate or desired. Some women, particularly those with no living children, mentioned pursuing a future pregnancy as a coping mechanism, or to enable them to regain social status. A few women took positivity from their experience, felt gratitude for their own survival, and were motivated to provide support for others in similar situations.

**Access to timely and culturally appropriate psychological support is valued.**(14,15,19–24,28,29,31,34,35,39,42–47) (FES 59%): Culturally and language appropriate psychological support, offered at an appropriate time, was effective in reducing anxiety, depression, and grief and may facilitate recovery for women. Women wanted healthcare workers to provide support, counselling, and good communication throughout their care. Access to appropriate support was limited however, by lack of trained or experienced staff, cost, and failure of referral by healthcare workers. The support needs of men may be overlooked.

**Women want information, advice and individualised discussions about future pregnancies.**(14,19–21,23–25,28,29,31,33,35,36,38–40,42,47) (FES 53%): Women reported mixed experiences of future pregnancy including fear of further loss, desire for another pregnancy and feeling pressure to conceive again soon. For some the thought of future pregnancy was helpful, although it was acknowledged that it would not be replacement for the loss. Women desired information, psychological support, and individualised discussions about future pregnancies.

**Addressing health system barriers is important for provision of respectful care.**(14,15,17,18,20,22–24,29,31,33,36,39,40,42–45) (FES 53%): Many women expressed dissatisfaction with the quality of care they received, including neglect, insensitivity, poor attitudes, and poor communication from healthcare workers. Staff, meanwhile, reported lack of sufficient resources, including facilities, equipment, and staff-shortages as barriers to providing good care. These factors contributed to delays in accessing care, along with lack of knowledge, female disempowerment and male or mother-in-law dominated decision making.

**Women may experience devaluation and stigmatization as a result of cultural practices and beliefs.**(14,22,23,27–33,36–38,40,41,43) (FES 47%): In the included LMIC studies, progressive social attitudes and empowerment of women was associated with ability to express and manage grief. Negative social perceptions of stillbirth, culture with male or mother-in-law decision making, expectation of a woman's role as child-bearer, son preference, and blaming women or curses/spirits for stillbirth instead of medical reasons, were associated with guilt, shame and domestic violence, and ultimately stigma for women. Tendency to suppression of mourning, lack of acknowledgement of motherhood after stillbirth, or absence of burial traditions may lead to disenfranchised grief

**Supporting proper investigation to understand causes of stillbirth may contribute to reducing stigma.**(20–23,27,29–31,33,36,38–40,44,45) (FES 44%): In the absence of a medical cause, women and communities relied on other explanations including superstitions and witchcraft, society, and poverty, often blaming themselves and others. Conversely, knowing the cause of death helped

women make sense of the loss and reduced fear of stigma. Women valued explanation of cause of stillbirth from healthcare workers, however, access to investigation of cause of death was limited by availability of autopsy and financial barriers.

**Women and staff believe that specialised bereavement care is important.**(15–

18,20,23,30,31,33,36,39,45) (*FES 35%*): Both women and staff described the need for specific guidelines to provide specialised care for women experiencing stillbirth. Suggestions included separation from women with livebirths, multidisciplinary and psychology input, and offering management choices including analgesia, appropriate to the local setting. Some staff were motivated to improve quality of care by their experience of poor outcomes, including stillbirth.

**Knowledge and information about stillbirth will empower women to take control of their own health.**(14,20,22–24,29,31,33,36,44,45,47) (*FES 35%*):

Women wanted more information, and opportunity for discussion, about delivery, cause of death, and postnatal care. In some settings this may also include postmortem and burial. This information, provided in their own language, would have helped to dispel fear of childbirth, fear for their own health, and given them control of their situation.

**Comprehensive staff training and support systems for staff are prerequisite to improving care.**(15–

18,20,24,45,47) (*FES 24%*): Both women and staff recognised that healthcare workers need more training in general communication and counselling skills, as well as specialist training in bereavement care and the needs of women after stillbirth. Staff caring for women with stillbirth also wanted support with the emotional impact, particularly in settings with high levels of perinatal loss where coping mechanisms and staff resilience were especially important.

**Women value supportive family presence throughout care.**(15,20,21,33,36,40,45) (*FES 21%*):

Family presence and involvement, especially from their spouse, from time of diagnosis and throughout care, was seen as beneficial by women. Women did not want to be left on their own.

**Women value follow-up care and advice to help them return to health.**(21,25,28,30,37) (*FES 15%*):

Both staff and patients considered continuity of care and follow up to be important. Women wanted advice on how to access further support after they had gone home, and on any underlying health problems. In LMICs stillbirth is more likely to be associated with physical morbidity or 'near-miss events' than in HICs, and women in this review reported long lasting sequelae, for both psychological and physical health.

**Intensity effect sizes:**

The intensity effect sizes for each study are included in appendix S5. Two studies contributed a larger proportion of findings(23,29), including after adjustment for findings with higher frequency effect sizes. These studies were both of medium quality and located in sub-Saharan Africa (Uganda and South Africa). The majority of the studies had an intensity effect size between 1.5 and 4.5, suggesting that a similar proportion of findings came from each study.



### **Quality assessment:**

Gough's weight of evidence (appendix S6) included study quality and risk of bias (rated low, medium or good), methodological and topic relevance (rated 1-3). No studies were of overall good quality, the majority (24/34) were of medium quality and rated 2 or 3 for relevance (21/34).

### **Discussion:**

#### **Main findings:**

This systematic review identified a range of literature addressing staff and parent experiences of care after stillbirth in LMICs. It also highlights the gaps in published literature on stillbirth experiences in many settings (Figure 1). There is considerable heterogeneity in cultural and healthcare provision among different LMICs and even within the same country. Further research needs to explore the generalisability of any findings.

Women experience a broad range of manifestations of grief following stillbirth across all settings, which may not be recognised by healthcare workers, or in their communities. This exacerbates negative experiences of stigmatisation, blame, devaluation, and loss of social status as a result of stillbirth. Positive attitudes and support during bereavement from family, communities and healthcare workers improves bereavement experience. Adequately developed health systems, with trained and supported staff, are better equipped to provide the support and information that women want after stillbirth.

#### **Strengths and Limitations:**

A key strength of this review is its inclusivity, involving a wide range of studies of varied methodology, population groups, and outcomes. The meta-summary method enables synthesis of relevant information from all different study types, whilst also quantifying the prevalence of each finding and the contribution of each paper to the whole. This enhances a comprehensive overview of the literature available.

The main limitation is the distribution of countries represented by the studies in this review (Figure 2). Gaps exist in areas with no research, whilst others may be over-represented by an individual researchers work; for example we found six papers for inclusion from one group in India(14,37,38,42,43,46). Cultural and healthcare practices may vary widely between, and even within, individual LMICs. Better understanding of the range of practices and experiences is needed before generalisations can be made. Deeply embedded cultural beliefs and practices related to stillbirth and bereavement mean that any findings must be interpreted with cultural sensitivity to any given location before implementation. Moreover, each individual woman's circumstances or preferences must be considered, before generalising findings to her.

The frequency effect sizes (FES) quoted should be interpreted as a description of prevalence in the literature, rather than clinical relevance or importance to women and families. A high FES may be influenced by multiple papers published by one research team with an interest in a particular topic, or conversely a low FES may reflect a lack of research rather than un-importance. As an illustration of this, in our review the theme around staff training and support has a low FES. This could be explained by only 4/34 studies explicitly addressing healthcare workers experiences, rather than indicating that the topic is un-important..

The individual studies involved in this review were mostly of low and medium quality. The meta-summary method considers that all studies may contribute useful information about the presence of themes despite issues of quality, and provision of intensity effect sizes (IES) (Appendix 5) allows readers to determine if any findings originate only from weaker studies. The description of study quality issues remains important, to drive improvements in quality in future studies. Some authors identified particular challenges including the richness of qualitative data improving in the absence of 'outsider' or 'western' investigators(27), and problems engaging staff with limited time for qualitative interviews(17) which could inform changes in future study design.

Finally, many of the principles and guidelines being proposed and investigated in other settings, and the themes and analysis following might still reflect Western sociocultural perspectives, particularly if the researchers were not native to the countries studied. It cannot therefore be assumed that findings can be imported to every LMIC, including the very countries the studies originated from, rather they would need to be critically adapted to local social norms and cultural practices first(30).

### **Interpretation:**

A key theme running through the findings was the negative experiences of women, particularly regarding, blame, stigmatisation, devaluation and lack of understanding among families, communities and healthcare workers, based on their reproductive status. This corresponds with a previous review of stillbirth outcomes which noted stigmatisation to be reported frequently in LMICs, compared to high-income countries (HICs)(6).

Improved awareness and public education about stillbirth could reduce the stigma experienced by women. It follows that increased understanding of the causes of stillbirth may reduce supernatural explanations in LMICs that focus blame on the woman. This correlates with recent findings about public perceptions of stillbirth in Ireland(48). In this high-income setting, it was found that the lack of awareness about causes of stillbirth led to blame of healthcare staff by the public. A common solution to this thread of misplaced blame may be public education, followed by appropriate stillbirth investigation, as allowed by local resources. Supporting investigation for causes of stillbirth, therefore, could help to reduce stigma in LMICs and HICs alike by focussing attention on biomedical explanations(49).

There is more emphasis on future pregnancy in the literature included in this review, than in the similar HIC review(5). This emphasis on future fertility in LMICs may come from a perception of women's value being linked to their reproductive status(25,27,29,41). Conversely, there was less emphasis on making memories, by seeing, holding and naming the baby, which may reflect cultural beliefs and practices.

The need for adequately equipped and developed health systems to provide care is clear in this review. Women report negative experiences related to poor attitudes and communication from health care workers, and healthcare workers report barriers to providing care including staff shortages and lack of training, facilities and equipment. Some interventions, such as developing localised guidelines, providing separate facilities for women experiencing stillbirth, and providing services for further investigation, would require financial and resource investment. Other changes, such as use of any available analgesia, and provision of respectful maternity care(50), require less material investment but could significantly change experience. Finally, some changes might not need investment, for example promoting support by family and friends during bereavement, but require changes in attitudes and behaviours that might not always be easy to achieve.

Overall, our findings correlate well with literature from HICs(5), suggesting that a set of common principles for bereavement care can be developed, and some of these principles could and should be promoted as a matter of urgency and without much resource.

Further research would be beneficial, particularly expanding the coverage of different cultures and countries to increase the breadth of the literature.

## **Conclusion:**

This systematic review shows that there is some literature available from LMICs providing insight into the experience of women, families and healthcare workers, with findings which could be used to inform improved care practices.

Further research is needed to better understand issues and design appropriate solutions, but there are already possible interventions that could make an immediate difference. These include public education to reduce stigma and blame and promoting the respectful maternity care agenda. The mere act of investigating stillbirth appropriately will send the signal to society that we must not blame women or staff. It is important to work with all stakeholders, clinicians, and politicians to implement improvements in bereavement care to reach all families. Educate, investigate, and respect, and we might be closer to improving global bereavement care than we think.

## **Acknowledgements:**

Disclosure of interests: Dimitrios Siassakos has received grants from SANDS (Stillbirth and Neonatal Death charity). No other authors have any conflicts of interests.

Contribution to authorship: DS conceived the study. DS, CSh and AM designed the study with the input of the wider group (CSt, HB, FB, VF, KG, DH, TM) . CSh, AM, DB, RB, KB performed searches, screening and data extraction. CSh, CSt, RB, ML, AM, DS analysed the data. CSh was responsible for writing the first draft and all authors contributed to finalising the manuscript.

Ethics approval: Not applicable

Funding: None

## **References:**

1. Heazell AEP, Siassakos D, Blencowe H, Burden C, Bhutta ZA, Cacciatore J, et al. Stillbirths: Economic and psychosocial consequences. *Lancet*. 2016;387(10018):604–16.
2. Lawn JE, Blencowe H, Waiswa P, Amouzou A, Mathers C, Hogan D, et al. Stillbirths : rates, risk factors, and acceleration towards 2030. *Lancet*. 2015;387:587–603.
3. Frederik Frøen J, Friberg IK, Lawn JE, Bhutta ZA, Pattinson RC, Allanson ER, et al. Stillbirths: Progress and unfinished business. *Lancet*. 2016;387(10018):574–86.
4. Bernis L De, Kinney M V, Stones W, Hooper-bender P, Vivio D, Leisher SH, et al. Stillbirths : ending preventable deaths by 2030. *Lancet*. 2016;387:703–16.

5. Ellis A, Chebsey C, Storey C, Bradley S, Jackson S, Flenady V, et al. Systematic review to understand and improve care after stillbirth : a review of parents ' and healthcare professionals ' experiences. *BMC Pregnancy Childbirth* [Internet]. BMC Pregnancy and Childbirth; 2016;16(16):1–19. Available from: <http://dx.doi.org/10.1186/s12884-016-0806-2>
6. Burden C, Bradley S, Storey C, Ellis A, Heazell AEP, Downe S, et al. From grief, guilt pain and stigma to hope and pride – a systematic review and meta-analysis of mixed-method research of the psychosocial impact of stillbirth. *BMC Pregnancy Childbirth* [Internet]. BMC Pregnancy and Childbirth; 2016;16(1):9. Available from: <http://www.biomedcentral.com/1471-2393/16/9>
7. World Bank Country and Lending Groups [Internet]. [cited 2018 Jan 11]. Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>
8. Covidence systematic review software. Melbourne, Australia: Veritas Health Innovation;
9. Vandenbroucke JP, von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): Explanation and Elaboration. *PLoS Med* [Internet]. 2007 Oct 16;4(10):e297. Available from: <http://dx.plos.org/10.1371/journal.pmed.0040297>
10. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Heal Care* [Internet]. 2007 Sep 16;19(6):349–57. Available from: <https://academic.oup.com/intqhc/article-lookup/doi/10.1093/intqhc/mzm042>
11. Gough D. Weight of evidence: a framework for the appraisal of the quality and relevance of evidence. *Appl Pract Res Ed Res Pap Educ*. 2007;22(2):213–28.
12. Sandelowski M, Barroso J, Voils C. Using Qualitative Metasummary to Synthesize Qualitative and Quantitative Descriptive Findings. *Res Nurs Heal*. 2007;30(1):99–111.
13. Sandelowski M, Barroso J. Creating Metasummaries of Qualitative Findings. *Nurs Res*. 2003;52(4):226–33.
14. Roberts LR, Montgomery SB, Ganesh G, Kaur HP, Singh R. Addressing Stillbirth in India Must Include Men. *Issues Ment Health Nurs*. 2017;1–10.
15. Santos C da S, Marques JF, Carvalho FHC, Fernandes AFC, Henriques ACPT, Moreira KDAP. Percepções de enfermeiras sobre a assistência prestada a mulheres diante do óbito fetal. *Esc Anna Nery*. 2012;16(2):277–84.
16. Moon Fai Chan, Lou F, Arthur DG. A Survey Comparing the Attitudes Toward Perinatal Bereavement Care of Nurses From Three Asian Cities. *Eval Health Prof* [Internet]. 2010;33(4):514–33. Available from: <http://journals.sagepub.com/doi/10.1177/0163278710381092>
17. Modiba L. Experiences and perceptions of midwives and doctors when caring for mothers with pregnancy loss in a Gauteng hospital. *Heal SA Gesondheid* [Internet]. 2008;13(4):29–40 12p. Available from: <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=105588562&site=ehost-live>
18. Petrites AD, Mullan P, Spangenberg K, Gold KJ. You have no choice but to go on: How physicians and midwives in Ghana cope with high rates of perinatal death. *Matern Child Health J*. 2016;20(7):1448–55.
19. Adeyemi A, Mosaku K, Ajenifuja O, Fatoye F, Makinde N, Ola B. Depressive Symptoms in a

- Sample of Women Following Perinatal Loss. *J Natl Med Assoc* [Internet]. Elsevier Masson SAS; 2008;100(12):1463–8. Available from: [http://dx.doi.org/10.1016/S0027-9684\(15\)31548-0](http://dx.doi.org/10.1016/S0027-9684(15)31548-0)
20. Allahdadian M, Irajpour A, Kazemi A, Kheirabadi G. Strategy for Mental Health Improvement of Iranian Stillborn Mothers From Their Perspective: A Qualitative Study. *Iran Red Crescent Med J* [Internet]. 2016;18(1):1–7. Available from: [http://www.ircmj.com/?page=article&article\\_id=21081](http://www.ircmj.com/?page=article&article_id=21081)
  21. Allahdadian M, Irajpour A, Kazemi A, Kheirabadi G. Social support: An approach to maintaining the health of women who have experienced stillbirth. *Iran J Nurs Midwifery Res* [Internet]. 2015;20(4):465–70. Available from: <http://www.ijnmrjournal.net/text.asp?2015/20/4/465/160998>
  22. Andajani-Sutjahjo S, Manderson L. Stillbirth, Neonatal Death and Reproductive Rights in Indonesia. *Reprod Health Matters* [Internet]. 2004 Jan 30;12(24):181–8. Available from: <https://www.tandfonline.com/doi/full/10.1016/S0968-8080%2804%2924141-6>
  23. Corbet-Owen C, Kruger L-M. The health system and emotional care: Validating the many meanings of spontaneous pregnancy loss. *Fam Syst Heal* [Internet]. 2001;19(4):411–27. Available from: <http://doi.apa.org/getdoi.cfm?doi=10.1037/h0089469>
  24. dos Santos ALD, Rosenberg CP, Buralli KO. [Histories of fetal losses told by women: research qualitative study]. *Rev Saude Publica* [Internet]. 2004 Apr;38(2):268–76. Available from: [http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S0034-89102004000200017&lng=pt&tIng=pt](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0034-89102004000200017&lng=pt&tIng=pt)
  25. Fottrell E, Kanhonou L, Goufodji S, Behague DP, Marshall T, Patel V, et al. Risk of psychological distress following severe obstetric complications in Benin: the role of economics, physical health and spousal abuse. *Br J Psychiatry* [Internet]. 2010 Jan 1;196(1):18–25. Available from: <http://bjp.rcpsych.org/cgi/doi/10.1192/bjp.bp.108.062489>
  26. Gausia K, Moran AC, Ali M, Ryder D, Fisher C, Koblinsky M. Psychological and social consequences among mothers suffering from perinatal loss: perspective from a low income country. *BMC Public Health* [Internet]. 2011 Dec 9;11(1):451. Available from: <http://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-11-451>
  27. Haws RA, Mashasi I, Mrisho M, Schellenberg JA, Darmstadt GL, Winch PJ. “These are not good things for other people to know”: How rural Tanzanian women’s experiences of pregnancy loss and early neonatal death may impact survey data quality. *Soc Sci Med* [Internet]. Elsevier Ltd; 2010;71(10):1764–72. Available from: <http://dx.doi.org/10.1016/j.socscimed.2010.03.051>
  28. Human M, Groenewald C, Odendaal HJ, Green S, Goldstein RD, Kinney HC. PSYCHOSOCIAL IMPLICATIONS OF STILLBIRTH FOR THE MOTHER AND HER FAMILY: A CRISIS-SUPPORT APPROACH. *Soc Work W* [Internet]. 2014 Nov 7;50(4):151–6. Available from: <http://socialwork.journals.ac.za/pub/article/view/392>
  29. Kiguli J, Namusoko S, Kerber K, Peterson S, Waiswa P. Weeping in silence: Community experiences of stillbirths in rural eastern Uganda. *Glob Health Action*. 2015;8(1):1–8.
  30. Kuti O, Ilesanmi CE. International Journal of Gynecology and Obstetrics Experiences and needs of Nigerian women after stillbirth. *Int J Gynecol Obstet* [Internet]. International Federation of Gynecology and Obstetrics; 2011;113(3):205–7. Available from: <http://dx.doi.org/10.1016/j.ijgo.2010.11.025>
  31. Modiba L, Nolte A. The experience of mothers who lost a baby during pregnancy. *Heal SA Gesondheid*. 2007;12(2):3–13.
  32. Obi SN, Onah HE, Okafor II. Depression among Nigerian women following pregnancy loss. *Int J*

- Gynecol Obstet [Internet]. International Federation of Gynecology and Obstetrics; 2009;105(1):60–2. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0020729208005675>
33. Osman HM, Egal JA, Kiruja J, Osman F, Byrskog U, Erlandsson K. Women's experiences of stillbirth in Somaliland: A phenomenological description. *Sex Reprod Healthc* [Internet]. Elsevier B.V.; 2017;11:107–11. Available from: <http://dx.doi.org/10.1016/j.srhc.2016.12.002>
  34. Paris GF, de Montigny F, Pelloso SM. Factors associated with the grief after stillbirth: A comparative study between Brazilian and Canadian women. *Rev da Esc Enferm*. 2016;50(4):546–53.
  35. Sutan R, Amin RM, Ariffin KB, Teng TZ, Kamal MF, Rusli RZ. Psychosocial impact of mothers with perinatal loss and its contributing factors: an insight. *J Zhejiang Univ Sci B* [Internet]. 2010;11(3):209–17. Available from: <http://www.springerlink.com/index/10.1631/jzus.B0900245>
  36. Sutan R, Miskam HM. Psychosocial impact of perinatal loss among Muslim women. 2012;
  37. Roberts LR, Montgomery S, Lee JW, Anderson BA. Social and Cultural Factors Associated with Perinatal Grief in Chhattisgarh, India. *J Community Health* [Internet]. 2012 Jun 29;37(3):572–82. Available from: <http://link.springer.com/10.1007/s10900-011-9485-0>
  38. Roberts LR, Anderson BA, Lee JW, Montgomery SB. Grief and Women: Stillbirth in the Social Context of India. *Int J Childbirth* [Internet]. 2012;2(3):187–98. Available from: <http://openurl.ingenta.com/content/xref?genre=article&issn=2156-5287&volume=2&issue=3&spage=187>
  39. Sereshti M, Nahidi F, Simbar M, Ahmadi F, Bakhtiari M, Zayeri F. Mothers' Perception of Quality of Services from Health Centers after Perinatal Loss. *Electron physician* [Internet]. 2016;8(2):2006–17. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4821318&tool=pmcentrez&rendertype=abstract>
  40. Sisay MM, Yirgu R, Gobezeayehu AG, Sibley LM. A qualitative study of attitudes and values surrounding stillbirth and neonatal mortality among grandmothers, mothers, and unmarried girls in rural Amhara and Oromiya regions, Ethiopia: Unheard souls in the backyard. *J Midwifery Women's Heal*. 2014;59(SUPPL1):S110–7.
  41. Attachie I., Mwini-Nyaledzigbor P, Affram C, Adjei D. Mothers' Experiences of Stillbirth : A Study in the Accra Metropolis ( Socio-Cultural Implications ) Mothers ' Experiences of Stillbirth : A Study in the Accra. *Adv Multidiscip Sci Res*. 2016;2(June):223–30.
  42. Roberts LR, Montgomery SB. Mindfulness-based Intervention for Perinatal Grief after Stillbirth in Rural India. *Issues Ment Health Nurs* [Internet]. 2015 Mar 4;36(3):222–30. Available from: <http://www.tandfonline.com/doi/full/10.3109/01612840.2014.962676>
  43. Roberts L, Montgomery S. Mindfulness-based Intervention for Perinatal Grief Education and Reduction among Poor Women in Chhattisgarh, India: a Pilot Study. *Interdiscip J best Pract Glob Dev* [Internet]. 2016 Apr;2(1):3047–54. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28357415>
  44. Simwaka ANK, de Kok B, Chilemba W. Women's perceptions of Nurse-Midwives' caring behaviours during perinatal loss in Lilongwe, Malawi: An exploratory study. *Malawi Med J*. 2014;26(1):8–11.
  45. Conry J, Prinsloo C. Mothers' access to supportive hospital services after the loss of a baby through stillbirth or neonatal death. *Heal SA Gesondheid*. 2008;13(2):14–24.

46. Roberts LR, Montgomery SB. Mindfulness-Based Intervention for Perinatal Grief in Rural India: Improved Mental Health at 12 Months Follow-Up. *Issues Ment Health Nurs*. 2016;37(12):942–51.
47. Chizhova MA. 127 PSYCHOLOGICAL HELP AFTER PERINATAL DEATH AS A PROPHYLACTIC OF REPRODUCTIVE PROBLEMS: CONTEMPORARY SITUATION IN RUSSIA. *Reprod Biomed Online* [Internet]. Reproductive Healthcare Ltd.; 2010 Oct;20:S54. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S147264831062545X>
48. Nuzum D, Meaney S, O'Donoghue K. The public awareness of stillbirth: an Irish population study. *BJOG An Int J Obstet Gynaecol* [Internet]. 2017 Oct 30; Available from: <http://doi.wiley.com/10.1111/1471-0528.14939>
49. Reinebrant H, Leisher S, Coory M, Henry S, Wojcieszek A, Gardener G, et al. Making stillbirths visible: a systematic review of globally reported causes of stillbirth. *BJOG An Int J Obstet Gynaecol* [Internet]. 2018 Jan;125(2):212–24. Available from: <http://doi.wiley.com/10.1111/1471-0528.14971>
50. WHO. The prevention and elimination of disrespect and abuse during facility-based childbirth [Internet]. [cited 2017 Oct 19]. Available from: [http://apps.who.int/iris/bitstream/10665/134588/1/WHO\\_RHR\\_14.23\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/134588/1/WHO_RHR_14.23_eng.pdf)

Figure 1: Prisma diagram

Figure 2: Study location and type

Figure 3: Thematic sentences

Figure 4: Frequency effect sizes

Appendix S1 Search strategy

Appendix S2 Data extraction form

Appendix S3 Summary of studies

Appendix S4 Sample extracted findings

Appendix S5 Intensity effect sizes

Appendix S6 Quality assessment

Appendix S7 GRIPP2-SF checklist